

Physics and Instrumentation of Atmospheric Aerosol Measurements Across Maryland July, 2011 **Communicating Scientific Research Methods to High School Physics Students** Ellen Babcock McLean, WT Woodson High School, Fairfax, VA NASA Mentors: M. Pippin, A. Omar

What are aerosols?


Aerosols are tiny particles of solids, liquids, or gases which become temporarily suspended in the air for varying lengths of time. Many particles, become airborne through natural means, but a significant portion of aerosols, especially the smaller ones, are anthropogenic: caused by Man.

In the atmosphere, aerosols can:

- Affect visibility
- behaviors in the atmosphere, and their affects human health.
- Form cloud condensation nuclei
- React with other particles forming hazardous compounds • Scatter, or absorb and then re-emit energy from photons
- Contribute to radiative forcing, an anthropogenic change in the radiation budget of the planet

In the biosphere aerosols can cause:

- Interference with the processes of plant respiration and photosynthesis
- cardiovascular disease
- Irregular heartbeat
- Aggravated asthma
- Decreased lung function
- Irritation of the airways
- Coughing
- Difficulty breathing



The size of aerosols can indicate

their source and potential

 PM_{10} are aerosols with diameters less than 10µm. Particles such as water, sand, pollen, soil, and sea salt are usually generated by natural events and tend to be classified as PM_{10} . They fall from the atmosphere after only a few hours or even minutes, giving them less opportunity to interact with the environment.

 $PM_{2.5}$ are aerosols with diameters less than 2.5 μ m. The majority of anthropogenic aerosols are classified as PM₂₅. Due to their small size, they can remain in the air for days, weeks or even months giving them increased opportunity to react with other substances in the atmosphere often with deleterious effect. PM_{25} are able to travel deep into the organs and tissues, remaining there for long periods of time and causing, or exacerbating, a host of health issues. PM₂₅'s longevity and potential for serious harm make it important to study and understand as, being largely anthropogenic, there are steps that can be taken towards positive change.

References

Readings to accompany the Physics of the Atmosphere, Physics Department East Stroudsburg University January 18, 11 Version 5.4 ©2011 by Robert A. Cohen retrieved 19 June, 2013 from p://www.esu.edu/physics/cohen/phys30 5/textbook/readings.5.4. Hanna, Steven R., et al. "Air Pollution." McGraw-Hill Encyclopedia of Science & Technology. 10th ed. Vol. 1. New York: McGraw-Hill, 2007. 292-309. Gale Virtual Reference Library. Web. 22 June 2013.

 Document URL http://go.galegroup.com/ps/i.do?id=GALE%7CCX3057500168&v=2.1&u=fair35939e&it=r&p=GVRL&sw=w •Zawar-Reza, Peyman. "Atmospheric Particulates Across Scales." Encyclopedia of Geography. Ed. Barney Warf. Vol. 1. SAGE Reference, 2010. 149-151. Gale Virtual Reference Library. Web. 22 June 2013.

 Document URL http://go.galegroup.com/ps/i.do?id=GALE%7CCX1788300073&v=2.1&u=fair35939e&it=r&p=GVRL&sw=w •Rollin, E. M. "An introduction to the use of Sun-photometry for the atmospheric

•correction of airborne sensor data." NERC EPFS Department of Geography University of Southampton Southampton, Electronic version retrieved from http://www.ncaveo.ac.uk/site-resources/pdf/cimel.pdf

 Holben, B. N., et al. AERONET—A Federated Instrument Network •Data Archive for Aerosol Characterization. REMOTE SENS. ENVIRON. 66:1-16 (1998) Retreived from

http://aeronet.gsfc.nasa.gov/new_web/PDF/afi.pdf

•BAM-1020 Operators Manual. From http://www.metone.com/aerosoldocs/BAM 3-sytem training.pdf 'Atmosphere." Encyclopedia of Public Health. Ed. Morton Lippmann, Lester Breslow. Vol. 1. New York: Macmillan Reference USA, 2002. 88-90. Gale Virtual Reference Library. Web. 22 June 2013. Document URL

http://go.galegroup.com/ps/i.do?id=GALE%7CCX3404000078&v=2.1&u=fair35939e&it=r&p=GVRL&sw=w "Aerosols." The Gale Encyclopedia of Science. 4th ed. Vol. 1. Detroit: Gale, 2008. 62-65. Gale Virtual Reference Library. Web. 22 June Document URL

http://go.galegroup.com/ps/i.do?id=GALE%7CCX2830100048&v=2.1&u=fair35939e&it=r&p=GVRL&sw=w •NASA Discover-AQ website http://www-air.larc.nasa.gov/missions/discover-aq/discover-aq.htm

•"Radiative Forcing." Environmental Science: In Context. Ed. Brenda Wilmoth Lerner and K. Lee Lerner. Vol. 2. Detroit: Gale, 2009. 680-682. In Context Series. Gale Virtual Reference Library. Web. 22 June 2013.

 Document URL http://go.galegroup.com/ps/i.do?id=GALE%7CCX3233900193&v=2.1&u=fair35939e&it=r&p=GVRL&sw=w •http://teaching.shu.ac.uk/hwb/chemistry/tutorials/molspec/beers1.htm

Acknowledgements

This work was supported by the Long-term Engagement in Authentic Research with NASA (LEARN) project with funding provided through a NASA SMD EPOESS grant.

Dr. Margaret Pippin for expertise, guidance, and inspiration, Dr. Ali Omar for his inside knowledge of aerosols and their physics, and Rebecca Kollmeyer for data retrieval and cleaning services


Aeronet and Discover AQ

Maryland EPA for sharing data

And the best last minute what equation do I use website ever: http://asdwww.larc.nasa.gov/SOLAR/experiment-index.html



Cimel-318 Automatic Sun Tracking Photometer Spectral extinction of direct beam radiation Aerosol optical thickness data is calculated by measuring voltage generated within a UV enhanced silicon detector of known surface area. 40-380-440-500-675-870-937 S1/S2 SUN




•The azimuth of the instrument adjusts automatically to keep the sun within the field of view (FOV in diagram above), and records the solar zenith angle, (Θ) (Fig2) •Optical air mass, (m), is calculated, (equation1) •Solar radiation passes through the first of two lenses, L₁(Fig1) • A disc housing 8 monochromatic filters is rotated in the path between the two lenses, L_1 and L_2 (Fig1) •The wavelength of light through each filter is recorded, (λ) • The silicon detector measures voltage generated by the

incident image, (v)

determined using data collected from reference instruments at NOAA's Mauna Loa Observatory in Hawaii Law (equation 2)

•Extraterrestrial voltage, (v_0) , is a calibration term •The Aerosol Optical Thickness is calculated using Beer's


m = --- $\sin\theta$ $\ln (v_{\lambda} - v_{0\lambda})$ (ea2)



 (F_{cal})

- measurement, (t)

- Be



Aeronet DRAGON (AOT)

FIG 3 AUTOMATIC SUNTRACKING SUNPHOTOMETER CE318_V2




Fig 1

 τ_{λ} = optical depth λ v = digital voltage v_0 = extraterrestrial voltage *m* = optical air mass = wavelength Θ = solar zenith angle

•Absorption per unit area is pre-determined during calibration

•Filter tape is exposed to radiation source

• A semi-conductor detector measures the intensity of beta ray transmission through the clean filter tape, (I_0)

•Air, at a controlled rate (V), is filtered through the tape allowing PM₂₅ to accumulate for the duration of the

•The semi-conductor detector measures the intensity of beta ray transmission through the dirty filter tape, (I) •The concentration of PM_{2.5} is calculated using the form of

10 **T**


 $C = concentration PM_{25}$ E_{cal} = calibration factor (mass density) V = measured mass flow rate t = measurement duration beta ray intensity at pre-exposure filter I = beta ray intensity at exposed filter

•The connection between measurement and data processing is the that connection provides the researcher with a far greater comprehension, and appreciation, of what that data represents.

•The quantities measured by remote sensing instrumentation are not the data reported by those instruments. Rather, the instrumentation has built into it mathematical models, or algorithms which translate the raw measurements into useable data.


•The Aeronet sun photometer column measurement and the EPA surface particulate monitor both provide indirect measurements of aerosol data each using different applications of Beer's Law


The amount of radiation absorbed or transmitted by a compound in a medium is proportional to the



•The graphed data suggests a strong correlation between the Aeronet and EPA measurements. The explanation of the importance of the data, the differences between what was actually measured, and the similarities between the data processing methods provides non or neo-scientists with more background information and so more understanding of the final product

can make that data more accessible to a wider range of viewers. •Making these tangible connections between science and the physical world can be used to teach the real scientific method to high school science students or even seize and hold the attention of a committed climate change skeptic.





3. Remote sensing data retrievals and current research methods will be included as curricular content in the physics classroom.

4. Students will learn valuable and marketable skill sets better preparing them for internships and their eventual undergraduate studies.